Acute liver failure or decompensated cirrhosis (4).

Calcii USP

rvastatin Tablets l

Removed 12/2022

Removed 12/2022

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ATORVASTATIN CALCIUM TABLETS safely and effectively. See full prescribing information for ATORVASTATIN CALCIUM TABLETS.

ATORVASTATIN CALCIUM TABLETS, for oral use

Initial U.S. Approval: 1996

RECENT MAJOR CHANGES-Contraindications, Pregnancy and Lactation (4)

Warnings and Precautions, CNS Toxicity (5.5) --INDICATIONS AND USAGE-

Atorvastatin calcium tablets is an HMG-CoA reductase inhibitor (statin) indicated (1): To reduce the risk of

- o Myocardial infarction (MI), stroke, revascularization procedures, and angina in adults with multiple risk factors for coronary heart disease (CHD) but without clinically evident
- o MI and stroke in adults with type 2 diabetes mellitus with multiple risk factors for CHD but without clinically evident CHD o Non-fatal MI, fatal and non-fatal stroke, revascularization procedures, hospitalization for congestive heart failure, and angina in adults with clinically evident CHD.
- As an adjunct to diet to reduce low-density lipoprotein (LDL-C) in: Adults with primary hyperlipidemia
- o Adults and pediatric patients aged 10 years and older with heterozygous familial hypercholesterolemia (HeFH).
- As an adjunct to other LDL-C-lowering therapies to reduce LDL-C in adults and pediatric patients aged 10 years and older with homozygous familial hypercholesterolemia.
- · As an adjunct to diet for the treatment of adults with
- Primary dysbetaliproteinemia Hypertriglyceridemia.

--DOSAGE AND ADMINISTRATION-

- Take orally once daily with or without food (2.1).
- Assess LDL-C when clinically appropriate, as early as 4 weeks after initiating Atorvastatin calcium tablets, and adjust dosage if necessary (2.1)
- o Recommended starting dosage is 10 or 20 mg once daily; dosage range is 10 mg to 80 mg once daily.
- o Patients requiring LDL-C reduction >45% may start at 40 mg once daily.
- Pediatric Patients Aged 10 Years of Age and Older with HeFH: Recommended starting dosage is 10 mg once daily; dosage range is 10 to 20 mg once daily (2.3).
- Pediatric Patients Aged 10 Years of Age and Older with HoFH: Recommended starting dosage is 10 to 20 mg once daily; dosage range is 10 to 80 mg once daily (2.4).

-DOSAGE FORMS AND STRENGTHS-

· See full prescribing information for Atorvastatin calcium tablets dosage modifications due to drug interactions (2.5).

Tablets: 10 mg, 20 mg, 40 mg, 80 mg of atorvastatin (3).

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

- 2 DOSAGE AND ADMINISTRATION
- Important Dosage Information
- 2.2 Recommended Dosage in Adult Patients
- Recommended Dosage in Pediatric Patients 10 Years of Age and Older with HeFH
- 2.4 Recommended Dosage in Pediatric Patients 10 Years of Age and Older with HoFH
- 2.5 Dosage Modifications Due to Drug Interactions

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Myopathy and Rhabdomyolysis
- Immune-Mediated Necrotizing Myopathy 5.3 Hepatic Dysfunction
- 5.4 Increases in HbA1c and Fasting Serum Glucose Levels
- 5.5 Increased Risk of Hemorrhagic Stroke in Patients on Atorvastatin calcium tablets 80 mg with Recent Hemorrhagic Stroke
- 6 ADVERSE REACTIONS

6.1 Clinical Trials Experience 6.2 Postmarketing Experience

- 7 DRUG INTERACTIONS $7.1 \quad \mathsf{Drug}\,\mathsf{Interactions}\,\mathsf{that}\,\mathsf{may}\,\mathsf{Increase}\,\mathsf{the}\,\mathsf{Risk}\,\mathsf{of}\,\mathsf{Myopathy}\,\mathsf{and}\,\mathsf{Rhabdomyolysis}\,\mathsf{with}\,\mathsf{atorvastatin}\,\mathsf{calcium}\,\mathsf{tablets}$
 - 7.2 Drug Interactions that may Decrease Exposure to atorvastatin calcium tablets 7.3 Atorvastatin calcium tablets Effects on Other Drugs

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy 8.2 Lactation

FULL PRESCRIBING INFORMATION

- 1 INDICATIONS AND USAGE . To reduce the risk of:
- $o\ \ Myocardial\ infarction\ (MI), stroke, revascularization\ procedures, and\ angina\ in\ adults\ with\ multiple\ risk\ factors$ for coronary heart disease (CHD) but without clinically evident CHD o MI and stroke in adults with type 2 diabetes mellitus with multiple risk factors for CHD but without clinically
- o Non-fatal MI, fatal and non-fatal stroke, revascularization procedures, hospitalization for congestive heart failure, and angina in adults with clinically evident CHD
- As an adjunct to diet to reduce low-density lipoprotein cholesterol (LDL-C) in:
- Adults with primary hyperlipidemia. $Adults\ and\ pediatric\ patients\ aged\ 10\ years\ and\ older\ with\ heterozygous\ familial\ hypercholesterolemia\ (HeFH).$
- $\bullet \quad \text{As an adjunct to other LDL-C-lowering therapies, or alone if such treatments are unavailable, to reduce LDL-C in}\\$ adults and pediatric patients aged 10 years and older with homozygous familial hypercholesterolemia (HoFH).
- · As an adjunct to diet for the treatment of adults with: o Primary dysbetalipoproteinemia
- o Hypertriglyceridemia
- 2 DOSAGE AND ADMINISTRATION 2.1 Important Dosage Information
- Take Atorvastatin calcium tablets orally once daily at any time of the day, with or without food. . Assess LDL-C when clinically appropriate, as early as 4 weeks after initiating Atorvastatin calcium tablets, and
- adjust the dosage if necessary. 2.2 Recommended Dosage in Adult Patients
- mmended starting dosage of Atorvastatin calcium tablets is 10 mg to 20 mg once daily. The dosage range is 10 mg to 80 mg once daily. Patients who require reduction in LDL-C greater than 45% may be started at 40 mg once
- 2.3 Recommended Dosage in Pediatric Patients 10 Years of Age and Older with HeFH The recommended starting dosage of Atorvastatin calcium tablets is 10 mg once daily. The dosage range is 10 mg to 20 mg once daily.
- 2.4 Recommended Dosage in Pediatric Patients 10 Years of Age and Older with HoFH ded starting dosage of Atorvastatin calcium tablets is 10 mg to 20 mg once daily. The dosage range is
- 10 mg to 80 mg once daily. 2.5 Dosage Modifications Due to Drug Interactions
- Concomitant use of Atorvastatin calcium tablets with the following drugs requires dosage modification of Atorvastatin calcium tablets [see Warnings and Precautions (5.1) and Drug Interactions (7.1)].
- In patients taking saguinavir plus ritonavir, darunavir plus ritonavir, fosamprenavir, fosamprenavir plus ritonavir.
- $elbas vir plus \ graz oprevir \ or \ letermovir, do \ not \ exceed \ Atorvas tatin \ calcium \ tablets \ 20 \ mg \ once \ daily.$ In patients taking nelfinavir, do not exceed Atorvastatin calcium tablets 40 mg once daily.
- Select Azole Antifungals or Macrolide Antibiotics
- $In patients \ taking \ clarithromycin \ or \ it raconazole, do \ not \ exceed \ Atorvas tatin \ calcium \ tablets \ 20 \ mg \ once \ daily.$ For additional recommendations regarding concomitant use of Atorvastatin calcium tablets with other anti-viral
- medications, azole antifungals or macrolide antibiotics, see Drug Interactions (7.1).

3 DOSAGE FORMS AND STRENGTHS Atorvastatin calcium tablets:

- 10 mg of atorvastatin: White to off-white, film-coated, oval shaped tablets "ATO" debossed on one side and "10" 20 mg of atorvastatin: White to off-white, film-coated, oval shaped tablets "ATO" debossed on one side and "20"
- 40 mg of atorvastatin: White to off-white, film-coated, oval shaped tablets "ATO" debossed on one side and "40"
- . 80 mg of atorvastatin: White to off-white, film-coated, oval shaped tablets "ATO" debossed on one side and "80 on other side.
- 4 CONTRAINDICATIONS
- Acute liver failure or decompensated cirrhosis [see Warnings and Precautions (5.3)]
 Hypersensitivity to atorvastatin or any excipients in atorvastatin calcium tablets. Hypersensitivity reactions including anaphylaxis, angioneurotic edema, erythema multiforme, Stevens-Johnson syndrome, and toxic epidermal necrolysis, have been reported [see Adverse Reactions (6.2)].
- 5 WARNINGS AND PRECAUTIONS

Atorvastatin calcium tablets may cause myopathy (muscle pain, tenderness, or weakness associated with elevated creatine kinase [CK]) and rhabdomyolysis. Acute kidney injury secondary to myoglobinuria and rare fatalities have occurred as a result of rhabdomyolysis in patients treated with statins, including atorvastatin calcium tablets.

Risk factors for myopathy include age 65 years or greater, uncontrolled hypothyroidism, renal impairment, concomitant use with certain other drugs (including other lipid-lowering therapies), and higher atorvastatin calcium tablets dosage [see Drug Interactions (7.1) and Use in Specific Populations (8.5, 8.6)].

Steps to Prevent or Reduce the Risk of Myopathy and Rhabdomyolysis

Atorvastatin calcium tablets exposure may be increased by drug interactions due to inhibition of cytochrome P450 enzyme 3A4 (CYP3A4) and/or transporters (e.g., breast cancer resistant protein [BCRP], organic anion-transporting polypeptide [OATP1B1/OATP1B3] and P-glycoprotein [P-gp]), resulting in an increased risk of myopathy and rhabdomyolysis. Concomitant use of cyclosporine, gemfibrozil, tipranavir plus ritonavir, or glecaprevir plus pibrentasvir with atorvastatin calcium tablets is not recommended. Atorvastatin calcium tablets dosage modifications are recommended for patients taking certain anti-viral, azole antifungals, or macrolide antibiotic medications [see Dosage and Administration (2.5)]. Cases of myopathy/rhabdomyolysis have been reported with atrovastatin co-administered with lipid modifying doses (>1 gram/day) of niacin, fibrates, colchicine, and ledipasvir plus sofosbuvir. Consider if the benefit of use of these products outweighs the increased risk of myopathy and rhabdomyolysis [see Drug Interactions (7.1)].

omitant intake of large quantities, more than 1.2 liters daily, of grapefruit juice is not recommended in patients taking atorvastatin calcium tablets [see Drug Interactions (7.1)].

Discontinue atorvastatin calcium tablets if markedly elevated CK levels occur or myopathy is either diagnosed or suspected. Muscle symptoms and CK elevations may resolve if atoryastatin calcium tablets is discontinued. Temporarily discontinue atorvastatin calcium tablets in patients experiencing an acute or serious condition at high risk of developing renal failure secondary to rhabdomyolysis (e.g., sepsis; shock; severe hypovolemia; major surgery; trauma; severe metabolic, endocrine, or electrolyte disorders; or uncontrolled epilepsy).

Inform patients of the risk of myopathy and rhabdomyolysis when starting or increasing the atorvastatin calcium tablets dosage. Instruct patients to promptly report any unexplained muscle pain, tenderness or weakness,

particularly if accompanied by malaise or fever

There have been rare reports of immune-mediated necrotizing myopathy (IMNM), an autoimmune myopathy associated with statin use, including reports of recurrence when the same or a different statin was administered. IMMM is characterized by proximal muscle weakness and elevated serum creatine kinase that persists despite discontinuation of statin treatment; positive anti-HMG CoA reductase antibody; muscle biopsy showing necrotizing myopathy; and improvement with immunosuppressive agents. Additional neuromuscular and serologic testing may be necessary. Treatment with immunosuppressive agents may be required. Discontinue atorvastatin calcium tablets if IMNM is suspected.

Increases in serum transaminases have been reported with use of atorvastatin calcium tablets [see Adverse Reactions (6.1)]. In most cases, these changes appeared soon after initiation, were transient, were not accompanied by symptoms, and resolved or improved on continued therapy or after a brief interruption in therapy. Persistent increases to more than three times the LILN in serum transaminases have occurred in approximately 0.7% of patients receiving astatin calcium tablets in clinical trials. There have been rare postmarketing reports of fatal and non-fatal hepatic failure in patients taking statins, including atorvastatin calcium tablets.

Patient who consume substantial quantities of alcohol and/or have a history of liver disease may be at increased risk 6.2 Postmarketing Experience for hepatic injury [see Use in Specific Populations (8.7)].

Consider liver enzyme testing before atorvastatin calcium tablets initiation and when clinically indicated thereafter. Atorvastatin calcium tablets is contraindicated in patients with acute liver failure or decompensated cirrhosis [see Contraindications (4)]. If serious hepatic injury with clinical symptoms and/or hyperbilirubinemia or jaundice occurs, promptly discontinue atoryastatin calcium tablets

5.4 Increases in HbA1c and Fasting Serum Glucose Levels Increases in HhA1c and fasting serum glucose levels have been reported with statins, including atoryastatin calcium

tablets. Optimize lifestyle measures, including regular exercise, maintaining a healthy body weight, and making

Hemorrhagic Stroke In a post-hoc analysis of the Stroke Prevention by Aggressive Reduction in Cholesterol Levels (SPARCL) trial where 2365 adult patients, without CHD who had a stroke or TIA within the preceding 6 months, were treated with atorvastatin calcium tablets 80 mg, a higher incidence of hemorrhagic stroke was seen in the atorvastatin calcium tablets 80 mg group compared to placebo (55, 2.3% atorvastatin calcium tablets vs. 33, 1.4% placebo; HR: 1.68, 95%

Increased Risk of Hemorrhagic Stroke in Patients on atorvastatin calcium tablets 80 mg with Recent

CI: 1.09, 2.59; p=0.0168). The incidence of fatal hemorrhagic stroke was similar across treatment groups (17 vs. 18 for the atorvastatin and placebo groups, respectively). The incidence of non-fatal hemorrhagic stroke was significantly higher in the atorvastatin calcium tablets group (38, 1.6%) as compared to the placebo group (16, 0.7%). Some baseline characteristics, including hemorrhagic and lacunar stroke on study entry, were associated with a higher incidence of hemorrhagic stroke in the atoryastatin calcium tablets group [see Adverse Reactions (6.1)]. Consider the risk/benefit of use of atorvastatin calcium tablets 80 mg in patients with recent hemorrhagic stroke ADVERSE REACTIONS

The following important adverse reactions are described below and elsewhere in the labeling:

- Myopathy and Rhabdomyolysis [see Warnings and Precautions (5.1)] Immune-Mediated Necrotizing Myopathy [see Warnings and Precautions (5.2)]
- Hepatic Dysfunction [see Warnings and Precautions (5.3)] Increases in HbA1c and Fasting Serum Glucose Levels [see Warnings and Precautions (5.4)]
- 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, the adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

In the atorvastatin calcium tablets placebo-controlled clinical trial database of 16,066 patients (8755 atorvastatin calcium tablets vs. 7311 placebo; age range 10–93 years, 39% women, 91% White, 3% Blacks, 2% Asians, 4% other) with a median treatment duration of 53 weeks, The most common adverse reactions in patients treated with atoryastatin calcium tablets that led to treatment discontinuation and occurred at a rate greater than placebo were: yalgia (0.7%), diarrhea (0.5%), nausea (0.4%), alanine aminotransferase increase (0.4%), and hepatic enzyme

 $Table \ 1 \ summarizes \ adverse \ reactions, \ reported \ in > 2\% \ and \ at \ a \ rate \ greater \ than \ placebo \ in \ patients \ treated \ with$ atorvastatin calcium tablets (n=8755), from seventeen placebo-controlled trials.

Table 1. Adverse Reactions Occurring in > 2% in Patients Atorvastatin calcium tablets-Treated with any Dose of and Greater than Placebo

Adverse Reaction	% Placebo N=7311	% 10 mg N=3908	% 20 mg N=188	% 40 mg N=604	% 80 mg N=4055	% Any dose N=8755
Managha and Wa						
Nasopharyngitis	8.2	12.9	5.3	7.0	4.2	8.3
Arthralgia	6.5	8.9	11.7	10.6	4.3	6.9
Diarrhea	6.3	7.3	6.4	14.1	5.2	6.8
Pain in extremity	5.9	8.5	3.7	9.3	3.1	6.0
Urinary tract infection	5.6	6.9	6.4	8.0	4.1	5.7
Dyspepsia	4.3	5.9	3.2	6.0	3.3	4.7
Nausea	3.5	3.7	3.7	7.1	3.8	4.0
Musculoskeletal pain	3.6	5.2	3.2	5.1	2.3	3.8
Muscle Spasms	3.0	4.6	4.8	5.1	2.4	3.6
Myalgia	3.1	3.6	5.9	8.4	2.7	3.5
Insomnia	2.9	2.8	1.1	5.3	2.8	3.0
Pharyngolaryngeal pain	2.1	3.9	1.6	2.8	0.7	2.3

Body as a whole: malaise, pyrexia

Digestive system: abdominal discomfort, eructation, flatulence, hepatitis, cholestasis Musculoskeletal system: musculoskeletal pain, muscle fatigue, neck pain, joint swelling

Metabolic and nutritional system: transaminases increase, liver function test abnormal, blood alkaline phosphatase Nervous system: nightmare Respiratory system: epistaxis Skin and appendages: urticaria

Urogenital system: white blood cells urine positive

atorvastatin group and (3.8%) of subjects in the placebo group

Special senses: vision blurred, tinnitus

Persistent elevations in serum transaminases, defined as more than 3 times the ULN and occurring on 2 or more occasions, occurred in 0.7% of patients who received atorvastatin calcium tablets in clinical trials. The incidence of these abnormalities was 0.2%, 0.2%, 0.6%, and 2.3% for 10, 20, 40, and 80 mg, respectively.

 $One \ patient \ in \ clinical \ trials \ developed \ jaun dice. \ Increases \ in \ liver \ enzyme \ tests \ in \ other \ patients \ were \ not \ associated$

with jaundice or other clinical signs or symptoms. Upon dose reduction, drug interruption, or discontinuation,

transaminase levels returned to or near pretreatment levels without seguelae. Eighteen of 30 patients with persistent liver enzyme elevations continued treatment with a reduced dose of atorvastatin calcium tablets In TNT [see Clinical Studies (14.1)] involving 10,001 subjects (age range 29-78 years, 19% women; 94% White, 3% Blacks 1% Asians 2% other) with clinically evident CHD treated with atoryastatin calcium tablets 10 mg daily (n=5006) or atorvastatin calcium tablets 80 mg daily (n=4995), in the high-dose atorvastatin calcium tablets group, there were more patients with serious adverse reactions (1.8%) and discontinuations due to adverse reactions (9.9%) as compared to the low-dose group (1.4%; 8.1%, respectively) during a median follow-up of 4.9 years. Persistent

and in 0.2% of individuals with atorvastatin 10 mg. Elevations of CK (> 10 x ULN) but were higher in the high-dose

atorvastatin group (0.3%) compared to the low-dose atorvastatin group (0.1%). $Stroke\ Prevention\ by\ Aggressive\ Reduction\ in\ Cholesterol\ Levels\ (SPARCL)$ In SPARCL 4731 patients (age range 21-92 years, 40% women; 93% White, 3% Black, 1% Asians, 3% other) without clinically evident CHD but with a stroke or transient ischemic attack (TIA) within the previous 6 months were treated with atorvastatin calcium tablets 80 mg (n=2365) or placebo (n=2366) for a median follow-up of 4.9 years, There was a higher incidence of persistent hepatic transaminase elevations (> 3 x ULN twice within 4-10 days) in the atorvastatin group (0.9%) compared to placebo (0.1%). Elevations of CK $(>10 \times ULN)$ were rare, but were higher in the atorvastatin group (0.1%) compared to placebo (0.0%). Diabetes was reported as an adverse reaction in (6.1%) of subjects in the

inase elevations (> 3 x ULN twice within 4–10 days) occurred in 1.3% of individuals with atoryastatin 80 mg

In a post-hoc analysis, atorvastatin calcium tablets 80 mg reduced the incidence of ischemic stroke (9.2% vs. 11.6%) and increased the incidence of hemorrhagic stroke (2.3% vs. 1.4%) compared to placebo. The incidence of fatal hemorrhagic stroke was similar between groups (17 atorvastatin calcium tablets vs. 18 placebo). The incidence of non-fatal hemorrhagic strokes was significantly greater in the atorvastatin group (38 non-fatal hemorrhagic strokes) as compared to the placebo group (16 non-fatal hemorrhagic strokes). Patients who entered the trial with a hemorrhagic stroke appeared to be at increased risk for hemorrhagic stroke (16% atorvastatin calcium tablets vs.

Adverse Reactions from Clinical Studies of atoryastatin calcium tablets in Pediatric Patients with HeFH

In a 26-week controlled study in *pediatric patients* with HeFH (ages 10 years to 17 years) (n=140, 31% female; 92% *White*, 1.6% Blacks, 1.6% Asians, 4.8% other), the safety and tolerability profile of atorvastatin calcium tablets 10 to 20 mg daily, as an adjunct to diet to reduce total cholesterol, LDL-C, and apo B levels, was generally similar to that of placebo [see Use in Specific Populations (8.4) and Clinical Studies (14.6)]. Hypersensitivity to atorvastatin or any excipient in Atorvastatin calcium tablets (4).

-WARNINGS AND PRECAUTIONS-

-CONTRAINDICATIONS

Myopathy and Rhabdomyolysis: Risk factors include age 65 years or greater, uncontrolled hypothyroidism, renal impairment, concomitant use with certain other drugs, and higher Atorvastatin calcium tablets dosage. Discontinue Atorvastatin calcium tablets if markedly elevated CK levels occur or myopathy is diagnosed or suspected. Temporarily discontinue Atorvastatin calcium tablets in patients experiencing an acute or serious condition at high risk of developing renal failure secondary to rhabdomyolysis. Inform patients of the risk of myopathy and rhabdomyolysis when starting or increasing Atorvastatin calcium tablets dosage. Instruct patients to promptly report unexplained muscle pain, tenderness, or weakness, particularly if accompanied by malaise or fever (2.5, 5.1, 7.1, 8.5, 8.6).

Immune-Mediated Necrotizing Myopathy (IMNM): Rare reports of IMNM, an autoimmune myopathy, have been reported with statin use. Discontinue Atorvastatin calcium tablets if IMNM is suspected (5.2).

Hepatic Dysfunction: Increases in serum transaminases have occurred, some persistent. Rare reports of fatal and non-fatal hepatic failure have occurred. Consider testing liver enzymes before initiating therapy and as clinically indicated thereafter. If serious hepatic injury with clinical symptoms and/or hyperbilirubinemia or jaundice occurs, promptly discontinue Atorvastatin calcium tablets (5.3).

-- ADVERSE REACTIONS-

Most common adverse reactions (incidence >5%) are nasopharyngitis, arthralgia, diarrhea, pain in extremity, and urinary tract infection (6.1).

To report SUSPECTED ADVERSE REACTIONS, contact www.umedicalabs.com at 1-855-288-577 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

-DRUG INTERACTIONS--

- See full prescribing information for details regarding concomitant use of Atorvastatin calcium tablets with other drugs or grapefruit juice that increase the risk of myopathy and rhabdomyolysis (2.5, 7.1).
- Rifampin: May reduce atorvastatin plasma concentrations. Administer simultaneously with Atorvastatin calcium tablets (7.2)
- Oral Contraceptives: May increase plasma levels of norethindrone and ethinyl estradiol; consider this effect when selecting an oral contraceptive (7.3).
- Digoxin: May increase digoxin plasma levels; monitor patients appropriately (7.3).
- -- USE IN SPECIFIC POPULATIONS--• Pregnancy: May cause fetal harm. (8.1).
- Lactation: Breastfeeding not recommended during treatment with Atorvastatin calcium tablets (8.2)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 10/2023

- 8.4 Pediatric Use
- 8.5 Geriatric Use
- Renal Impairment 8.6 8.7 Hepatic Impairment
- 10 OVERDOSAGE
- 11 DESCRIPTION
- 12 CLINICAL PHARMACOLOGY 12.1 Mechanism of Action
- 12.2 Pharmacodynamics 12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 14 CLINICAL STUDIES 16 HOW SUPPLIED/STORAGE AND HANDLING
- 17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed

The following adverse reactions have been identified during post-approval use of atorvastatin calcium tablets. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate reventing or managing them

their frequency or establish a causal relationship to drug exposure. Gastrointestinal disorders: pancreatitis

General disorders: fatique Hepatobiliary Disorders: fatal and non-fatal hepatic failure

Respiratory disorders: interstitial lung disease

Immune system disorders: anaphylaxis

Injury: tendon rupture Musculoskeletal and connective tissue disorders: rhabdomyolysis, myositis

 $There \ have \ been \ rare \ reports \ of \ immune-mediated \ necrotizing \ myopathy \ associated \ with \ statin \ use.$ There have been rare reports of cognitive impairment (e.g., memory loss, forgetfulness, amnesia, memory impairment, confusion) associated with the use of all statins. Cognitive impairment was generally nonserious, and

reversible upon statin discontinuation, with variable times to symptom onset (1 day to years) and symptom resolution

Skin and subcutaneous tissue disorders: angioneurotic edema, bullous rashes (including erythema multiforme, Stevens-Johnson syndrome, and toxic epidermal necrolysis)

DRUG INTERACTIONS 7.1 Drug Interactions that may Increase the Risk of Myopathy and Rhabdomyolysis with atorvastatin calcium Atorvastatin calcium tablets is a substrate of CYP3A4 and transporters (e.g., OATP1B1/1B3, P-gp, or BCRP) Atorvastatin calcium tablets plasma levels can be significantly increased with concomitant administration of inhibitors of CYP3A4 and transporters. Table 2 includes a list of drugs that may increase exposure to atorvastatin calcium tablets

and may increase the risk of myopathy and rhabdomyolysis when used concomitantly and instructions for preventing or managing them [see Warnings and Precautions (5.1) and Clinical Pharmacology (12.3)]. Table 2: Drug Interactions that may Increase the Risk of Myopathy and Rhabdomyolysis with atorvastatin calcium

lets	
closporine or Gemfibrozil	
Clinical Impact:	Atorvastatin plasma levels were significantly increased with concomitant administration of atorvastatin calcium tablets and cyclosporine, an inhibitor of CYP3A4 and OATP1B1[see Clinical Pharmacology (12.3)]. Gemfibrozil may cause myopathy when given alone. The risk of myopathy and rhabdomyolysis is increased with concomitant use of cyclosporine or gemfibrozil with atorvastatin calcium tablets.
Intervention:	Concomitant use of cyclosporine or gemfibrozil with atorvastatin calcium tablets are not recommended.
ti-Viral Medications	
	Atorvastatin plasma levels were significantly increased with concomitant

are inhibitors of CYP3A4 and/or transporters (e.g., BCRP, OATP1B1/1B3, P-gp, MRP2, and/or OAT2) [see Clinical Pharmacology (12.3)]. Cases of myopathy and habdomyolysis have been reported with concomitant use of ledipasvir plus ofosbuvir with atorvastatin calcium tablets. Concomitant use of tipranavir plus ritonavir or glecaprevir plus pibrentasvir with atorvastatin calcium tablets is not recommended.

In patients taking lopinavir plus ritonavir, or simeprevir, consider the risk/benefit of concomitant use with atorvastatin. In patients taking saquinavir plus ritonavir, darunavir plus ritonavir, fosampren fosamprenavir plus ritonavir, elbasvir plus grazoprevir or letermovir, do not exceed atorvastatin calcium tablets 20 mg.

In patients taking nelfinavir, do not exceed atorvastatin calcium tablets 40 mg [see Dosage and Administration (2.5)1 Consider the risk/benefit of concomitant use of ledipasvir plus sofosbuvir with atorvastatin calcium tablets. Monitor all patients for signs and symptoms of myopathy particularly during initiation of therapy and during upward dose titration of either drug. Tipranavir plus ritonavir, glecaprevir plus pibrentasvir, lopinavir plus ritonavir

Select Azole Antifungals or Macrolide Antibiotics orvastatin plasma levels were significantly increased with concon dministration of atorvastatin calcium tablets with select azole antifungals or Pharmacology (12.3)]. In patients taking clarithromycin or itraconazole, do not exceed atorvastatin calcii tablets 20 mg [see Dosage and Administration (2.5)]. Consider the risk/benefit of

rythromycin, clarithromycin, itraconazole, ketoconazole, posaconazole, and Fxamnle Cases of myopathy and rhabdomyolysis have been observed with concomitant use of Clinical Impact: lipid modifying dosages of niacin (>1 gram/day niacin) with atorvastatin calciu nsider if the benefit of using lipid modifying dosages of niacin concom atorvastatin calcium tablets outweighs the increased risk of myopathy and habdomyolysis. If concomitant use is decided, monitor patients for signs and symptoms of myopathy particularly during initiation of therapy and during upward

tablets outweighs the increased risk of myopathy and rhabdomyolysis. If concomitant use is decided, monitor patients for signs and symptoms of myopathy particularly during initiation of therapy and during upward dose titration of either drug. Cases of myopathy and rhabdomyolysis have been reported with concomitant use o olchicine with atorvastatin calcium tablets. Consider the risk/benefit of concomitant use of colchicine with atorvastatin calc

Grapefruit juice consumption, especially excessive consumption, more than Clinical Impact: 1.2 liters/daily, can raise the plasma levels of atorvastatin and may increase the risk of yopathy and rhabdomyolysis. void intake of large quantities of grapefruit juice, more than 1.2 liters daily, who taking atorvastatin calcium tablets.

7.2 Drug Interactions that may Decrease Exposure to atorvastatin calcium tablets Table 3 presents drug interactions that may decrease exposure to atorvastatin calcium tablets and instructions for

Pharmacology (12.3)].

of cytochrome P450 3A4 and inhibitor of OATP1B1, can lead to variable reductions in plasma concentrations of atorvastatin. Due to the dual interaction mechanism of Clinical Impact: rifampin, delayed administration of atorvastatin calcium tablets after administration of

Table 3: Drug Interactions that may Decrease Exposure to atorvastatin calcium tablets

rifampin has been associated with a significant reduction in atorvastatin plasma Intervention: Administer atorvastatin calcium tablets and rifampin simultaneously. 7.3 Atorvastatin calcium tablets Effects on Other Drugs

Table 4 presents atorvastatin calcium tablets effect on other drugs and instructions for preventing or managing them

Oral Contraceptives Co-administration of atoryastatin calcium tablets and an oral contracentive increased Clinical Impact: plasma concentrations of norethindrone and ethinyl estradiol [see Clinical Pharmacology (12.3)]. Intervention. calcium tablets. Digoxin When multiple doses of atorvastatin calcium tablets and digoxin were co-

Clinical Impact: administered, steady state plasma digoxin concentrations increased [see Clinical

Intervention: Monitor patients taking digoxin appropriately USE IN SPECIFIC POPULATIONS

Discontinue atorvastatin calcium tablets when pregnancy is recognized. Alternatively, consider the ongoing therapeutic needs of the individual patient. Atorvastatin calcium tablets decreases synthesis of cholesterol and possibly other biologically active substances derived from cholesterol; therefore, atorvastatin calcium tablets may cause fetal harm when administered to pregnant patients based on the mechanism of action [see Clinical Pharmacology (12.71)]. In addition, treatment of hyperlipidemia is not generally necessary during pregnancy. Atherosclerosis is a chronic process and the discontinuation of lipid-lowering drugs during pregnancy should have little impact on the outcome of long-term therapy of primary hyperlipidemia for most patients. Available data from case series and prospective and retrospective observational cohort studies over decades of use

reproduction studies, no adverse developmental effects were observed in pregnant rats or rabbits orally administered atorvastatin at doses that resulted in up to 30 and 20 times, respectively, the human exposure at the maximum recommended human dose (MRHD) of 80 mg, based on body surface area (mg/m2). In rats administered atorvastatin during gestation and lactation, decreased postnatal growth and development delay were observed at doses \geq 6 times The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

A Medicaid cohort linkage study of 1152 statin-exposed pregnant women compared to 886,996 controls did not find a

with statins in pregnant women have not identified a drug-associated risk of major congenital malformations. Published data from prospective and retrospective observational cohort studies with atorvastatin calcium tablets use in pregnant women are insufficient to determine if there is a drug-associated risk of miscarriage (see Data). In animal

significant teratogenic effect from maternal use of statins in the first trimester of pregnancy, after adjusting for potential confounders – including maternal age, diabetes mellitus, hypertension, obesity, and alcohol and tobacco use using propensity score-based methods. The relative risk of congenital malformations between the group with static use and the group with no statin use in the first trimester was 1.07 (95% confidence interval 0.85 to 1.37) afte use and the group with no stain use in the inst trimester was 1.07 (95% conholence interval 0.86 to 1.37) after controlling for confounders, particularly pre-existing diabetes mellitus. There were also no statistically significant increases in any of the organ-specific malformations assessed after accounting for confounders. In the majority of pregnancies, statin treatment was initiated prior to pregnancy and was discontinued at some point in the first trimester when pregnancy was identified. Study limitations include reliance on physician coding to define the presence of a malformation, lack of control for certain confounders such as body mass index, use of prescription dispensing as verification for the use of a statin, and lack of information on non-live births.

Animal Data Atomastatin was administered to pregnant rats and rabbits during organogenesis at oral doses up to 300 mg/kg/day and 100 mg/kg/day, respectively. Atomastatin was not teratogenic in rats at doses up to 300 mg/kg/day or in rabbits at doses up to 100 mg/kg/day. These doses resulted in multiples of about 30 times (rat) or 20 times (rabbit) the human exposure at the MRHD based on surface area (mg/m2). In rats, the maternally toxic dose of 300 mg/kg resulted in increased post-implantation loss and decreased fetal body weight. At the maternally toxic doses of 50 and 100 mg/kg/day in rabbits, there was increased post-implantation loss, and at 100 mg/kg/day fetal body weights were

In a study in pregnant rats administered 20, 100, or 225 mg/kg/day from gestation day 7 through to lactation day 20 (weaning), there was decreased survival at birth, postnatal day 4, weaning, and post-weaning in pups of mothers dosed with 225 mg/kg/day, a dose at which maternal toxicity was observed. Pup body weight was decreased through postnatal day 21 at 100 mg/kg/day, and through postnatal day 91 at 225 mg/kg/day. Pup development was delayed rotarod performance at 100 mg/kg/day and acoustic startle at 225 mg/kg/day; pinnae detachment and eye-opening at 225 mg/kg/day). These doses correspond to 6 times (100 mg/kg) and 22 times (225 mg/kg) the human exposure at the MBND based at MC.

8.2 Lactation There is no information about the presence of atorvastatin in human milk, the effects of the drug on the breastfed infan or the effects of the drug on milk production. However, it has been shown that another drug in this class passes into human milk. Studies in rats have shown that atorvastatin and/or its metabolites are present in the breast milk of lactating rats. When a drug is present in animal milk, it is likely that the drug will be present in human milk (see Data).

Atorvastatin crosses the rat placenta and reaches a level in fetal liver equivalent to that of maternal plasma.

Statins, including atorvastatin calcium tablets, decrease cholesterol synthesis and possibly the synthesis of other biologically active substances derived from cholesterol and may cause harm to the breastfed infant. Because of the potential for serious adverse reactions in a breastfed infant, based on the mechanism of action, advise atients that breastfeeding is not recommended during treatment with atorvastatin calcium tablets [see Use in Specific Populations (8.1), Clinical Pharmacology (12.1)].

Following a single oral administration of 10 mg/kg of radioactive atorvastatin to lactating rats, the concentration of total radioactivity was determined. Atorvastatin and/or its metabolites were measured in the breast milk and pup plasma at a 2:1 ratio (milk:plasma). The safety and effectiveness of atorvastatin calcium tablets as an adjunct to diet to reduce LDL-C have been established pediatric patients 10 years of age and older with HeFH. Use of atorvastatin calcium tablets for this indication is based on a double-blind, placebo-controlled clinical trial in 187 pediatric patients 10 years of age and older with HeFH. In this limited controlled trial, there was no significant effect on growth or sexual maturation in the boys or girls, or on

The safety and effectiveness of atorvastatin calcium tablets as an adjunct to other LDL-C-lowering therapies to reduce LDL-C have been established pediatric patients 10 years of age and older with HoFH. Use of atorvastatin calcium tablets for this indication is based on a trial without a concurrent control group in 8 pediatric patients 10 years of age and older with HoFH [see Clinical Studies (14)].

The safety and effectiveness of atorvastatin calcium tablets have not been established in pediatric patients younger than 10 years of age with HeFH or HoFH, or in pediatric patients with other types of hyperlipidemia (other than HeFH or

menstrual cycle length in girls.

400.00 mm

simeprevir, saquinavir plus ritonavir, darunavir plus ritonavir, fosamprenavir, fosamprenavir plus ritonavir, elbasvir plus grazoprevir, letermovir, nelfinavir, and

oncomitant use of other azole antifungals or macrolide antibiotics with atorvastatin calcium tablets. Monitor all patients for signs and symptoms of myopathy particularly luring initiation of therapy and during upward dose titration of either drug.

e titration of either drug. Fibrates may cause myopathy when given alone. The risk of myopathy and habdomyolysis is increased with concomitant use of fibrates with atorvastating onsider if the benefit of using fibrates concomitantly with atorvastatin calcium

tablets. If concomitant use is decided, monitor patients for signs and symptoms of yopathy particularly during initiation of therapy and during upward dose titration of

8.5 Geriatric Use Of the total number of atorvastatin calcium tablets -treated patients in clinical trials, 15.813 (40%) were >>65 years old and 2.800 (7%) were > 75 years old. No overall differences in safety or effectiveness were observed between these

Advanced age ($\geq \geq 65$ years) is a risk factor for atorvastatin calcium tablets -associated myopathy and rhabdomyolysis. Dose selection for an elderly patient should be cautious, recognizing the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy and the higher risk of myopathy. Monitor geriatric patients receiving atorvastatin calcium tablets for the increased risk of myopathy [see Warnings and Precautions (5.1) and Clinical Pharmacology (12.3)].

Renal impairment is a risk factor for myopathy and rhabdomyolysis. Monitor all patients with renal impairment for development of myopathy. Renal impairment does not affect the plasma concentrations of atorvastatin calcium tablets, therefore there is no dosage adjustment in patients with renal impairment *[see Warnings and Precautions (5.1)*

In patients with chronic alcoholic liver disease, plasma concentrations of atorvastatin calcium tablets are markedly increased. Cmax and AUC are each 4-fold greater in patients with Childs-Pugh A disease. Cmax and AUC are approximately 16-fold and 11-fold increased, respectively, in patients with Childs-Pugh B disease. Atorvastatin calcium tablets is contraindicated in patients with acute liver failure or decompensated cirrhosis [see Contraindications (4)1.

10 OVERDOSAGE $No\ specific\ antidotes\ for\ Atorvastatin\ calcium\ tablets\ are\ known.\ Contact\ Poison\ Control\ (1-800-222-1222)\ for\ latest$ recommendations. Due to extensive drug binding to plasma proteins, hemodialysis is not expected to significantly

11 DESCRIPTION

enhance atorvastatin calcium clearance.

Atorvastatin is an inhibitor of 3-hydroxy-3-methylglutaryl-coenzyme A (HMG-CoA) reductase.

Atorvastatin calcium is [R-R*, R*)]-2-(4-fluorophenyl)-β, δ-dihydroxy-5-(1-methylethyl)-3-phenyl-4-[(phenylamino) $carbonyl] \hbox{-1H-pyrrole-1-heptanoic acid, calcium salt (2:1) trihydrate. The empirical formula of atorva statin calcium is a continuous carbonyl] \hbox{-1H-pyrrole-1-heptanoic acid, calcium salt (2:1) trihydrate.} \\$ (C33H34 FN2O5)2Ca•3H2O and its molecular weight is 1209.42. Its structural formula is:

Atorvastatin calcium is a white to off-white powder that is insoluble in aqueous solutions of pH 4 and below Atorvastatin calcium is very slightly soluble in distilled water, pH 7.4 phosphate buffer, and acetonitrile; slightly soluble in ethanol; and freely soluble in methanol.

orvastatin calcium tablets, USP for oral use contain atorvastatin 10 mg, 20 mg, 40 mg, or 80 mg (equivalent to 10.359 mg, 20.718 mg, 41.436 mg, or 82.872 mg atorvastatin calcium trihydrate) and the following inactive ingredients: Croscarmellose sodium, NF; Hydroxy propyl cellulose, NF; Lactose monohydrate, NF; Magnesium stearate, NF; Microcrystalline cellulose, NF; Polysorbate 80, NF; Precipitated calcium carbonate, NF; Opadry Complete $film\, coating\, system\, YS\text{--}1\text{--}7040\, White\, (Hypromellose,\, Macrogol,\, Talc\, and\, Titanium\, dioxide).$

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Atorvastatin calcium is a selective, competitive inhibitor of HMG-CoA reductase, the rate-limiting enzyme that converts 3-hydroxy-3-methylglutaryl-coenzyme A to mevalonate, a precursor of sterols, including cholesterol. In animal models, atorvastatin calcium lowers plasma cholesterol and lipoprotein levels by inhibiting HMG-CoA reductase and cholesterol synthesis in the liver and by increasing the number of hepatic LDL receptors on the cell surface to enhance uptake and catabolism of LDL; atorvastatin calcium also reduces LDL production and the number of LDL particles.

Atorvastatin calcium, as well as some of its metabolites, are pharmacologically active in humans. The liver is the primary site of action and the principal site of cholesterol synthesis and LDL clearance. Drug dosage, rather than systemic drug concentration, correlates better with LDL-C reduction. Individualization of drug dosage should be based on therapeutic response [see Dosage and Administration (2)].

Atorvastatin calcium tablets are rapidly absorbed after oral administration; maximum plasma concentrations occur within 1 to 2 hours. Extent of absorption increases in proportion to atorvastatin calcium tablets dose. The absolute bioavailability of atorvastatin (parent drug) is approximately 14% and the systemic availability of HMG-CoA reductase inhibitory activity is approximately 30%. The low systemic availability is attributed to presystemic clearance in gastrointestinal mucosa and/or hepatic first-pass metabolism. Although food decreases the rate and extent of drug absorption by approximately 25% and 9%, respectively, as assessed by Cmax and AUC, LDL-C reduction is similar whether atorvastatin calcium tablets are given with or without food. Plasma atorvastatin calcium concentrations are lower (approximately 30% for Cmax and AUC) following evening drug administration compared with morning. However, LDL-C reduction is the same regardless of the time of day of drug administration.

 $Mean \ volume \ of \ distribution \ of \ atorva statin \ calcium \ is \ approximately \ 381 \ liters. \ Atorva statin \ calcium \ is \ > 98\% \ bound$ to plasma proteins. A blood/plasma ratio of approximately 0.25 indicates poor drug penetration into red blood cells.

Atorvastatin calcium is extensively metabolized to ortho- and parahydroxylated derivatives and various beta-oxidation products. In vitro inhibition of HMG-CoA reductase by ortho- and parahydroxylated metabolites is equivalent to that of atorvastatin calcium. Approximately 70% of circulating inhibitory activity for HMG-CoA reductase is attributed to active metabolites. In vitro studies suggest the importance of atoryastatin calcium metabolism by cytochrome P450 3A4, consistent with increased plasma concentrations of atorvastatin calcium in humans following co-administration with erythromycin, a known inhibitor of this isozyme [see Drug Interactions (7.1)]. In animals, the ortho-hydroxy metabolite undergoes further glucuronidation.

Atorvastatin calcium and its metabolites are eliminated primarily in bile following hepatic and/or extra-hepatic metabolism; however, the drug does not appear to undergo enterohepatic recirculation. Mean plasma elimination halflife of atorvastatin calcium in humans is approximately 14 hours, but the half-life of inhibitory activity for HMG-CoA reductase is 20 to 30 hours due to the contribution of active metabolites. Less than 2% of a dose of atorvastatin calcium is recovered in urine following oral administration.

Specific Populations

entrations of atorvastatin calcium are higher (approximately 40% for Cmax and 30% for AUC) in healthy There were 61 deaths in the atorvastatin calcium tablets group vs. 82 deaths in the placebo group (HR 0.73, p=0.059). elderly subjects (age >65 years) than in young adults.

Apparent oral clearance of atorvastatin in pediatric subjects appeared similar to that of adults when scaled allometrically by body weight as the body weight was the only significant covariate in atorvastatin population PK model with data including pediatric HeFH patients (ages 10 years to 17 years of age, n=29) in an open-label, 8-week study.

Plasma concentrations of atorvastatin calcium tablets in women differ from those in men (approximately 20% higher for Cmax and 10% lower for AUC); however, there is no clinically significant difference in LDL-C reduction with

 $Renal\ disease\ has\ no\ influence\ on\ the\ plasma\ concentrations\ or\ LDL-C\ reduction\ of\ atorvastatin\ calcium\ [see\ Use\ in\ Plasma\ concentrations\ or\ LDL-C\ reduction\ of\ atorvastatin\ calcium\ [see\ Use\ in\ Plasma\ concentration\ or\ LDL-C\ reduction\ of\ atorvastatin\ calcium\ [see\ Use\ in\ Plasma\ concentration\ or\ LDL-C\ reduction\ of\ atorvastatin\ calcium\ [see\ Use\ in\ Plasma\ concentration\ or\ LDL-C\ reduction\ of\ atorvastatin\ calcium\ [see\ Use\ in\ Plasma\ concentration\ or\ LDL-C\ reduction\ of\ atorvastatin\ calcium\ [see\ Use\ in\ Plasma\ concentration\ or\ LDL-C\ reduction\ of\ atorvastatin\ calcium\ [see\ Use\ in\ Plasma\ concentration\ or\ LDL-C\ reduction\ of\ atorvastatin\ calcium\ [see\ Use\ in\ Plasma\ concentration\ or\ LDL-C\ reduction\ of\ atorvastatin\ calcium\ [see\ Use\ in\ Plasma\ concentration\ or\ LDL-C\ reduction\ of\ atorvastatin\ calcium\ [see\ Use\ in\ Plasma\ concentration\ or\ LDL-C\ reduction\ of\ atorvastatin\ calcium\ [see\ Use\ in\ Use\ concentration\ concentration\ or\ concentration$

Specific Populations (8.6)]. While studies have not been conducted in patients with end-stage renal disease, hemodialysis is not expected to significantly enhance clearance of Atorvastatin calcium since the drug is extensively bound to plasma pro

In patients with chronic alcoholic liver disease, plasma concentrations of atorvastatin calcium are markedly increased Cmax and AUC are each 4-fold greater in patients with Childs-Pugh A disease. Cmax and AUC are approximately 16-fold and 11-fold increased, respectively, in patients with Childs-Pugh B disease [see Use in Specific Populations (8.7)].

Attorvastatin is a substrate of the hepatic transporters, OATP1B1 and OATP1B3 transporter. Metabolites of atorvastatin In the Treating to New Targets Study (TNT), the effect of atorvastatin calcium tablets 80 mg/day vs. atorvastatin are substrates of OATP1B1. Atorvastatin is also identified as a substrate of the efflux transporter BCRP, which may limit the intestinal absorption and biliary clearance of atorvastatin.

Table 5: Effect of Co-administered Drugs on the Pharmacokinetics of Atorvastatin

Co-administered drug and dosing regimen	A	torvastatin	
	Dose (mg)	Ratio of AUC&	Ratio of Cmax 8
#Cyclosporine 5.2 mg/kg/day,stable dose	10 mg QDa for 28 days	8.69	10.66
#Tipranavir 500 mg BIDb/ritonavir 200 mg BIDb, 7 days	10 mg, SDc	9.36	8.58
#Glecaprevir 400 mg QDa/pibrentasvir 120 mg QDa, 7 days	10 mg QDa for 7 days	8.28	22.00
#Telaprevir 750 mg q8hf, 10 days	20 mg, SDc	7.88	10.60
#; Saquinavir 400 mg BIDb/ ritonavir 400 mg BIDb, 15 days	40 mg QDa for 4 days	3.93	4.31
#Elbasvir 50 mg QDa/grazoprevir 200 mg QDa, 13 days	10 mg SDc	1.94	4.34
#Simeprevir 150 mg QDa, 10 days	40 mg SDc	2.12	1.70
#Clarithromycin 500 mg BIDb, 9 days	80 mg QDa for 8 days	4.54	5.38
#Darunavir 300 mg BIDb/ritonavir 100 mg BIDb, 9 days	10 mg QDa for 4 days	3.45	2.25
#Itraconazole 200 mg QDa, 4 days	40 mg SDc	3.32	1.20
#Letermovir 480 mg QDa, 10 days	20 mg SDc	3.29	2.17
#Fosamprenavir 700 mg BIDb/ritonavir 100 mg BIDb, 14 days	10 mg QDa for 4 days	2.53	2.84
#Fosamprenavir 1400 mg BIDb, 14 days	10 mg QDa for 4 days	2.30	4.04
#Nelfinavir 1250 mg BIDb, 14 days	10 mg QDa for 28 days	1.74	2.22
#Grapefruit Juice, 240 mL QDa,*	40 mg SDc	1.37	1.16
Diltiazem 240 mg QDa, 28 days	40 mg SDc	1.51	1.00
Erythromycin 500 mg QIDe, 7 days	10 mg SDc	1.33	1.38
Amlodipine 10 mg, single dose	80 mg SDc	1.18	0.91
Cimetidine 300 mg QIDe, 2 weeks	10 mg QDa for 2 weeks	1.00	0.89
Colestipol 10 g BIDb, 24 weeks	40 mg QDa for 8 weeks	NA	0.74**
MaaloxTC® 30 mL QIDe, 17 days	10 mg QDa for 15 days	0.66	0.67
Efavirenz 600 mg QDa, 14 days	10 mg for 3 days	0.59	1.01
#Rifampin 600 mg QDa, 7 days (co-administered) [†]	40 mg SDc	1.12	2.90
#Rifampin 600 mg QDa, 5 days (doses separated) [†]	40 mg SDc	0.20	0.60
#Gemfibrozil 600 mg BIDb, 7 days	40 mg SDc	1.35	1.00
#Fenofibrate 160 mg QDa, 7 days	40 mg SDc	1.03	1.02
Boceprevir 800 mg TIDd, 7 days	40 mg SDc	2.32	2.66

- $Represents\ ratio\ of\ treatments\ (co-administered\ drug\ plus\ atorvastatin\ vs.\ atorvastatin\ alone).$ See Sections 5.1 and 7 for clinical significance.
- Greater increases in AUC (ratio of AUC up to 2.5) and/or Cmax (ratio of Cmax up to 1.71) have been reported with excessive grapefruit consumption (>750 mL 1.2 liters per day). Ratio based on a single sample taken 8-16 h post dose.
- recommended, as delayed administration of atorvastatin after administration of rifampin has been associated with a significant reduction in atorvastatin plasma concentrations
- The dose of saquinavir plus ritonavir in this study is not the clinically used dose. The increase in atorvastati exposure when used clinically is likely to be higher than what was observed in this study. Therefore, caution should be applied and the lowest dose necessary should be used.
- Once daily Twice daily
- Single dose Three times daily

Four times daily Every 8 hours

Table 6: Effect of Atorvastatin on the Pharmacokinetics of Co-administered Drugs						
Atorvastatin	rvastatin Co-administered drug and dosing regimen					
	Drug/Dose (mg)	Ratio of AUC	Ratio of Cmax			
80 mg QDa for 15 days	Antipyrine, 600 mg SDc	1.03	0.89			
80 mg QDa for 10 days	# Digoxin 0.25 mg QDa, 20 days	1.15	1.20			
40 mg QDa for 22 days	Oral contraceptive QDa, 2 months - norethindrone 1 mg - ethinyl estradiol 35µg	1.28 1.19	1.23 1.30			
10 mg, SDc	Tipranavir 500 mg BIDb/ritonavir 200 mg BIDb, 7 days	1.08	0.96			
10 mg QDa for 4 days	Fosamprenavir 1400 mg BIDb, 14 days	0.73	0.82			
10 mg QDa for 4 days	Fosamprenavir 700 mg	0.99	0.94			

- # See Section 7 for clinical significance. Once daily
- Twice daily c Single dose

Atorvastatin calcium tablets had no clinically significant effect on prothrombin time when administered to patients

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In a 2-year carcinogenicity study in rats at dose levels of 10, 30, and 100 mg/kg/day, 2 rare tumors were found in muscle in high-dose females; in one, there was a rhabdomyosarcoma and, in another, there was a fibrosarcoma. This ats a plasma AUC (0-24) value of approximately 16 times the mean human plasma drug exposure after an

A 2-year carcinogenicity study in mice given 100, 200, or 400 mg/kg/day resulted in a significant increase in liver adenomas in high-dose males and liver carcinomas in high-dose females. These findings occurred at plasma AUC (0-24) values of approximately 6 times the mean human plasma drug exposure after an 80 mg oral dose

In vitro, atorvastatin was not mutagenic or clastogenic in the following tests with and without metabolic activation: the Ames test with Salmonella typhimurium and Escherichia coli, the HGPRT forward mutation assay in Chinese hamster In female rats, atorvastatin at doses up to 225 mg/kg (56 times the human exposure) did not cause adverse effects on fertility. Studies in male rats performed at doses up to 175 mg/kg (15 times the human exposure) produced no changes in fertility. There was aplasia and aspermia in the epididymis of 2 of 10 rats treated with 100 mg/kg/day of atorvastatin for 3 months (16 times the human AUC at the 80 mg dose); testis weights were significantly lower at 20 and 100 mg/kg/day of add 100 mg/kg and colificantly lower at 20 and 100 mg/kg/day of atorvastatin rational significantly lower at 20 and 100 mg/kg/day of atorvastatin rational significantly lower at 20 and 100 mg/kg/day of atorvastatin rational significantly lower at 20 and 100 mg/kg/day of atorvastatin rational significantly lower at 20 and 100 mg/kg/day of atorvastatin rational significantly lower at 20 and 100 mg/kg/day of atorvastatin rational significantly lower at 20 and 100 mg/kg/day of atorvastatin rational significantly lower at 20 and 100 mg/kg/day of atorvastatin rational significantly lower at 20 and 100 mg/kg/day of atorvastatin rational significantly lower at 20 and 100 mg/kg/day of atorvastatin rational significantly lower at 20 at 100 mg/kg/day of atorvastatin rational significantly lower at 20 at 100 mg/kg/day of atorvastatin rational significantly lower at 20 at 100 mg/kg/day of atorvastatin rational significantly lower at 20 at 100 mg/kg/day of atorvastatin rational significantly lower at 20 at 100 mg/kg/day of atorvastatin rational significantly lower at 20 at 100 mg/kg/day of atorvastatin rational significantly lower at 20 at 100 mg/kg/day of atorvastatin rational significantly lower at 20 at 100 mg/kg/day of atorvastatin rational significantly lower at 20 at 100 mg/kg/day of atorvastatin rational significantly lower at 20 at 100 mg/kg/day of atorvastatin rational significantly lower at 20 at 100 mg/kg/day of atorvastatin rational significantly lower at 20 at 100 mg/kg/day of 20 atorvastatin rational significantly lower at 20 at 100 mg/kg/d

and 100 mg/kg and epididymal weight was lower at 100 mg/kg. Male rats given 100 mg/kg/day for 11 weeks prior to mating had decreased sperm motility, spermatid head concentration, and increased abnormal sperm. Atorvastatin

caused no adverse effects on semen parameters, or reproductive organ histopathology in dogs given doses of 10, 40,


or 120 mg/kg for 2 years.

14 CLINICAL STUDIES

Prevention of Cardiovascular Disease In the Anglo-Scandinavian Cardiac Outcomes Trial (ASCOT), the effect of atorvastatin calcium tablets on fatal and non-fatal coronary heart disease was assessed in 10,305 patients with hypertension, 40–80 years of age (mean of 63 years, and other coronary heart disease was assessed in 10,305 patients with hypertension, 40–80 years of age (mean of 63 years, and other coronary heart disease was assessed in 10,305 patients with hypertension, 40–80 years of age (mean of 63 years, and other coronary heart disease was assessed in 10,305 patients with hypertension, 40–80 years of age (mean of 63 years, and other coronary heart disease was assessed in 10,305 patients with hypertension, 40–80 years of age (mean of 63 years, and other coronary heart disease was assessed in 10,305 patients with hypertension, 40–80 years of age (mean of 63 years, and other coronary heart disease was assessed in 10,305 patients with hypertension, 40–80 years of age (mean of 63 years, and 04 years). 19% women; 95% White, 3% Black, 1% South Asian, 1% other), without a previous myocardial infarction and with total cholesterol (TC) levels <251 mg/dL. Additionally, all patients had at least 3 of the following cardiovascular risk factors: male gender (81%), age >55 years (85%), smoking (33%), diabetes (24%), history of CHD in a first-degree relative (26%),TC: HDL >6 (14%), peripheral vascular disease (5%), left ventricular hypertrophy (14%), prior cerebrovascular event (10%), specific ECG abnormality (14%), proteinuria/albuminuria (62%). In this double-blind placebo-controlled trial, patients were treated with anti-hypertensive therapy (goal BP 140/90 mm Hg for non-diabetic patients; <130/80 mm Hg for patients with diabetes) and allocated to either atorvastatin calcium tablets 10 mg daily (n=5168) or placebo (n=5137), using a covariate adaptive method which took into account the distribution of nine baseline characteristics of patients already enrolled and minimized the imbalance of those characteristics across the groups. Patients were followed for a median duration of 3.3 years.

The effect of 10 mg/day of atorvastatin calcium tablets on lipid levels was similar to that seen in previous clinical trials. Atorvastatin calcium tablets significantly reduced the rate of coronary events [either fatal coronary heart disease (46 events in the placebo group vs. 40 events in the atorvastatin calcium tablets group) or non-fatal MI (108 events in the placebo group vs. 60 events in the atorvastatin calcium tablets group) with a relative risk reduction of 36% [(based on incidences of 1.9% for atorvastatin calcium tablets vs. 3.0% for placebo), p=0.0005 (see Figure 1)]. The risk reduction was consistent regardless of age, smoking status, obesity, or presence of renal dysfunction. The effect of atorvastatin calcium tablets was seen regardless of baseline LDL levels.

Figure 1: Effect of atorvastatin calcium tablets 10 mg/day on Cumulative Incidence of Non-Fatal Myocardial tion or Coronary Heart Disease Death (in ASCOT-LLA)



Atorvastatin calcium tablets also significantly decreased the relative risk for revascularization procedures by 42% (incidences of 1.4% for atorvastatin calcium tablets and 2.5% for placebo). Although the reduction of fatal and non-fatal strokes did not reach a pre-defined significance level (p=0.01), a favorable trend was observed with a 26% relative risk reduction (incidences of 1.7% for atorvastatin calcium tablets and 2.3% for placebo). There was no significant difference between the treatment groups for death due to cardiovascular causes (p=0.51) or noncardio


In the Collaborative Atorvastatin Diabetes Study (CARDS), the effect of atorvastatin calcium tablets on cardiovascular disease (CVD) endpoints was assessed in 2838 subjects (94% white, 2% Black, 2% South Asian, 1% other; 68% male), ages 40–75 with type 2 diabetes based on WHO criteria, without prior history of cardiovascular disease and with LDL < 160 mg/dL and triglycerides (TG) < 600 mg/dL. In addition to diabetes, subjects had 1 or more of the following risk factors: current smoking (23%), hypertension (80%), retinopathy (30%), or microalbuminuria (9%) or macroalbuminuria (3%). No subjects on hemodialysis were enrolled in the trial. In this multicenter, placebo controlled, double-blind clinical trial, subjects were randomly allocated to either atorvastatin calcium tablets10 mg daily (1429) or placebo (1411) in a 1:1 ratio and were followed for a median duration of 3.9 years. The primary endpoint was the occurrence of any of the major cardiovascular events; myocardial infarction, acute CHD death, unstable angina, coronary revascularization, or stroke. The primary analysis was the time to first occurrence of the primary

 $Base line\ characteristics\ of\ subjects\ were:\ mean\ age\ of\ 62\ years,\ mean\ HbA1c\ 7.7\%;\ median\ LDL-C\ 120\ mg/dL;\ median\ age\ of\ 62\ years,\ mean\ HbA1c\ 7.7\%;\ median\ LDL-C\ 120\ mg/dL;\ median\ LDL-C\ 120\ mg/$ TC 207 mg/dL; median TG 151 mg/dL; median HDL-C 52 mg/dL

The effect of atorvastatin calcium tablets 10 mg/day on lipid levels was similar to that seen in previous clinical trials. Atorvastatin calcium tablets significantly reduced the rate of major cardiovascular events (primary endpoint events) (83 events in the atorvastatin calcium tablets group vs. 127 events in the placebo group) with a relative risk reduction of 37%, HR 0.63, 95% CI (0.48, 0.83) (p=0.001) (see Figure 2). An effect of atorvastatin calcium tablets was seen regardless of age, sex, or baseline lipid levels.

Atorvastatin calcium tablets significantly reduced the risk of stroke by 48% (21 events in the atorvastatin calcium tablets group vs. 39 events in the placebo group), HR 0.52, 95% CI (0.31, 0.89) (p=0.016) and reduced the risk of MI by 42% (38 events in the atorvastatin calcium tablets group vs. 64 events in the placebo group), HR 0.58, 95.1% CI (0.39, 0.86) (p=0.007). There was no significant difference between the treatment groups for angina, revascularization

Figure 2: Effect of atorvastatin calcium tablets 10 mg/day on Time to Occurrence of Major Cardiovascular Event ocardial infarction, acute CHD death, unstable angina, coronary revascularization, or stroke) in CARDS



calcium tablets 10 mg/day on the reduction in cardiovascular events was assessed in 10.001 subjects (94% white. 81% male, 38% > 65 years) with clinically evident coronary heart disease who had achieved a target LDL-Clevel<130 mg/dL after completing an 8-week, open-label, run-in period with atorvastatin calcium tablets 10 mg/day. Subjects were randomly assigned to either 10 mg/day or 80 mg/day of atorvastatin calcium tablets and followed for a median duration of 4.9 years. The primary endpoint was the time-to-first occurrence of any of the following major cardiovascular events (MCVE): death due to CHD, non-fatal myocardial infarction, resuscitated cardiac arrest, and fatal and non-fatal stroke. The mean LDL-C, TC, TG, non-HDL, and HDL cholesterol levels at 12 weeks were 73, 145, 128, 98, and 47 mg/dL during treatment with 80 mg of atorvastatin calcium tablets and 99, 177, 152, 129, and 48 mg/dL ment with 10 mg of atorvastatin calcium tablets.

Treatment with atorvastatin calcium tablets 80 mg/day significantly reduced the rate of MCVE (434 events in the 80 mg/day group vs. 548 events in the 10 mg/day group) with a relative risk reduction of 22%, HR 0.78, 95% CI (0.69, 0.89), p=0.0002 (see Figure 3 and Table 7). The overall risk reduction was consistent regardless of age (<65, >65) or

Figure 3: Effect of atorvastatin calcium tablets 80 mg/day vs. 10 mg/day on Time to Occurrence of Major Cardiovascular Events (TNT)




Table 7: Overview of Efficacy Results in TNT						
Endpoint	Atorvastatin Endpoint 10 mg (N=5006)		80	astatin mg 1995)	HRa (95%CI)	
PRIMARY ENDPOINT	n	(%)	n	(%)		
First major cardiovascular endpoint	548	(10.9)	434	(8.7)	0.78 (0.69, 0.89)	
Components of the Primary Endpoint						
CHD death	127	(2.5)	101	(2.0)	0.80 (0.61, 1.03)	
Non-fatal, non-procedure related MI	308	(6.2)	243	(4.9)	0.78 (0.66, 0.93)	
Resuscitated cardiac arrest	26	(0.5)	25	(0.5)	0.96 (0.56, 1.67)	
Stroke (fatal and non-fatal)	155	(3.1)	117	(2.3)	0.75 (0.59, 0.96)	
SECONDARY ENDPOINTS*						
First CHF with hospitalization	164	(3.3)	122	(2.4)	0.74 (0.59, 0.94)	
First PVD endpoint	282	(5.6)	275	(5.5)	0.97 (0.83, 1.15)	
First CABG or other coronary	904	(18.1)	667	(13.4)	0.72 (0.65, 0.80)	
revascularization procedureb						
First documented angina endpointb	615	(12.3)	545	(10.9)	0.88 (0.79, 0.99)	
All-cause mortality	282	(5.6)	284	(5.7)	1.01 (0.85, 1.19)	
Components of All-Cause Mortality						
Cardiovascular death	155	(3.1)	126	(2.5)	0.81 (0.64, 1.03)	
Noncardiovascular death	127	(2.5)	158	(3.2)	1.25 (0.99, 1.57)	
Cancer death	75	(1.5)	85	(1.7)	1.13 (0.83, 1.55)	
Other non-CV death	43	(0.9)	58	(1.2)	1.35 (0.91, 2.00)	
Suicide, homicide, and othertraumatic	9	(0.2)	15	(0.3)	1.67 (0.73, 3.82)	
non-CV death		1				

^a Atorvastatin 80 mg: atorvastatin 10 mg Component of other secondary endpoints

* Secondary endpoints not included in primary endpoint

HR=hazard ratio; CHD=coronary heart disease; CI=confidence interval; MI=myocardial infarction; CHF=congestive heart failure; CV=cardiovascular; PVD=peripheral vascular disease; CABG=coronary artery bypass graft Confidence intervals for the Secondary Endpoints were not adjusted for multiple comparisons

Of the events that comprised the primary efficacy endpoint, treatment with atorvastatin calcium tablets 80 mg/day significantly reduced the rate of non-fatal, non-procedure related MI and fatal and non-fatal stroke, but not CHD death or resuscitated cardiac arrest (Table 7). Of the predefined secondary endpoints, treatment with atorvastatin calcium tablets 80 mg/day significantly reduced the rate of coronary revascularization, angina, and hospitalization for heart failure, but not peripheral vascular disease. The reduction in the rate of CHF with hospitalization was only observed in the 8% of patients with a prior history of CHF.

There was no significant difference between the treatment groups for all-cause mortality (Table 7). The proportions of subjects who experienced cardiovascular death, including the components of CHD death and fatal stroke, were numerically smaller in the atorvastatin calcium tablets 80 mg group than in the atorvastatin calcium tablets 10 mg treatment group. The proportions of subjects who experienced noncardiovascular death were numerically larger in the atorvastatin calcium tablets 80 mg group than in the atorvastatin calcium tablets 10 mg treatment group.

Primary Hyperlipidemia in Adults

Atorvastatin calcium tablets reduces total-C, LDL-C, apo B, and TG, and increases HDL-C in patients with hyperlipidemia (heterozygous familial and nonfamilial) and mixed dyslipidemia. Therapeutic response is seen within 2 weeks, and maximum response is usually achieved within 4 weeks and maintained during chronic therapy.

In two multicenter, placebo-controlled, dose-response trials in patients with hyperlipidemia, atorvastatin calcium tablets given as a single dose over 6 weeks, significantly reduced total-C, LDL-C, apo B, and TG. (Pooled

able 8: Dose Response in Patients With Primary Hyperlipidemia (Adjusted Mean % Change From Baseline)						
Dose	N	TC	LDL-C	Apo B	TG	HDL-C
Placebo	21	4	4	3	10	-3
10	22	-29	-39	-32	-19	6
20	20	-33	-43	-35	-26	9
40	21	-37	-50	-42	-29	6
80	23	-45	-60	-50	-37	5

Treatment (Daily Dose)	N	Total-C	LDL-C	Apo B	TG	HDL-C
Trial 1						
atorvastatin calcium	707	-27a	-36a	-28a	-17a	+7
tablets 10 mg						
Lovastatin 20 mg	191	-19	-27	-20	-6	+7
95% CI for Diff ¹		-9.2, -6.5	-10.7, -7.1	-10.0, -6.5	-15.2, -7.1	-1.7, 2.0
Trial 2						
atorvastatin calcium	222	-25b	-35b	-27b	-17b	+6
tablets 10 mg						
Pravastatin 20 mg	77	-17	-23	-17	-9	+8
95% CI for Diff ¹		-10.8, -6.1	-14.5, -8.2	-13.4,-7.4	-14.1, -0.7	-4.9, 1.6
Trial 3						
atorvastatin calcium	132	-29c	-37c	-34c	-23c	+7
tablets 10 mg						
Simvastatin 10 mg	45	-24	-30	-30	-15	+7
95% CI for Diff1		-8.72.7	-10.12.6	-8.01.1	-15.10.7	-4.3. 3.9

 1 A negative value for the 95% CI for the difference between treatments favors atorvastatin calcium tablets for all except HDL-C, for which a positive value favors atoryastatin calcium tablets. If the range does not include 0, this indicates a statistically significant difference.

Significantly different from lovastatin, ANCOVA, p < 0.05

Significantly different from pravastatin, ANCOVA, p < 0.05 Significantly different from simvastatin, ANCOVA, p < 0.05

Table 9 does not contain data comparing the effects of atorvastatin calcium tablets 10 mg and higher doses of lovastatin, pravastatin, and simvastatin. The drugs compared in the trials summarized in the table are not necessarily interchangeable.

The response to atorvastatin calcium tablets in 64 patients with isolated hypertriglyceridemia treated across several clinical trials is shown in the table below (Table 10). For the atorvastatin calcium-treated patients, median (min, max)

Table 10: Combined Patients With Isolated Elevated TG: Median (min, max) Percentage Change from Baseline

	Placebo (N=12)	atorvastatin calcium tablets 10 mg (N=37)	atorvastatin calcium tablets 20 mg (N=13)	atorvastatin calcium tablets 80 mg (N=14)
Triglycerides	-12.4 (-36.6, 82.7)	-41.0 (-76.2, 49.4)	-38.7 (-62.7, 29.5)	-51.8 (-82.8, 41.3)
Total-C	-2.3 (-15.5, 24.4)	-28.2 (-44.9, -6.8)	-34.9 (-49.6, -15.2)	-44.4 (-63.5, -3.8)
LDL-C	3.6 (-31.3, 31.6)	-26.5 (-57.7, 9.8)	-30.4 (-53.9, 0.3)	-40.5 (-60.6, -13.8)
HDL-C	3.8 (-18.6, 13.4)	13.8 (-9.7, 61.5)	11.0 (-3.2, 25.2)	7.5 (-10.8, 37.2)
non-HDL-C	-2.8 (-17.6, 30.0)	-33.0 (-52.1, -13.3)	-42.7 (-53.7, -17.4)	-51.5 (-72.9, -4.3)

<u>Dysbetalipoproteinemia in Adults</u>

The results of an open-label crossover trial of 16 patients (genotypes: 14 apo E2/E2 and 2 apo E3/E2) with dysbetalipoproteinemia are shown in the table below (Table 11).

Table 11: Onen-Label Crossover Trial of 16 Patients With Dysheta

Tubio I I. Opon Lubo	TOTOGOGOVOT THAT OF TOT ALTOHIO W	iii byobotanpoprotomom	ia (i rourionoon typo iii)
		Median	% Change (min, max)
	Median (min, max) at Baseline (mg/dL)	Atorvastatin calcium tablets 10 mg	Atorvastatin calcium tablets 80 mg
Total-C	442 (225, 1320)	-37 (-85, 17)	-58 (-90, -31)
Triglycerides	678 (273, 5990)	-39 (-92, -8)	-53 (-95, -30)
IDL-C + VLDL-C	215 (111, 613)	-32 (-76, 9)	-63 (-90, -8)
non-HDL-C	411 (218, 1272)	-43 (-87, -19)	-64 (-92, -36)

HoFH in Adults and Pediatric Patients

In a trial without a concurrent control group, 29 patients (mean age of 22 years, median age of 24 years, (31% <18 years) with HoFH received maximum daily doses of 20 to 80 mg of atorvastatin calcium tablets. The mean LDL-C reduction in this trial was 18%. Twenty-five patients with a reduction in LDL-C had a mean response of 20% (range of 7% to 53%, median of 24%); the remaining 4 patients had 7% to 24% increases in LDL-C. Five of the 29 patients had absent LDL-receptor function. Of these, 2 patients also had a portacaval shunt and had no significant reduction in

$LDL-C.\ The\ remaining\ 3\ receptor-negative\ patients\ had\ a\ mean\ LDL-C\ reduction\ of\ 22\%.$

In a double-blind, placebo-controlled study followed by an open-label phase, 187 boys and post-menarchal girls 10 years to 17 years of age (mean age 14.1 years; 31% female; 92% White, 1.6% Blacks, 1.6% Asians, 4.8% other) with heterozygous familial hypercholesterolemia (HeFH) or severe hypercholesterolemia, were randomized to atorvastatin calcium tablets (n=140) or placebo (n=47) for 26 weeks and then all received atorvastatin calcium tablets for 26 weeks. Inclusion in the trial required 1) a baseline LDL-C level > 190 mg/dL and positive family history of FH or documented premature cardiovascular disease in a first or second-degree relative. The mean baseline LDL-C value was 219mg/dL (range: 139-385 mg/dL) in the atorvastatin calcium tablets group compared to 230 mg/dL (range: 160-325 mg/dL) in the placebo group. The dosage of atorvastatin calcium tablets (once daily) was 10 mg for the first 4 weeks and uptitrated to 20 mg if the LDL-C level was > 130 mg/dL. The number of -treated patients who required uptitration to 20 mg after Week 4 during the double-blind phase was 78

Atorvastatin calcium tablets significantly decreased plasma levels of total-C, LDL-C, triglycerides, and apolipoprotein B during the 26-week double-blind phase (see Table 12).

How do I store atorvastatin calcium tablets?

• Store atorvastatin calcium tablets at nome

Table 12: Lipid-altering Effects of Atorvastatin Calcium Tablets in Adolescent Boys and Girls with Heterozygou Familial Hypercholesterolemia or Severe Hypercholesterolemia (Mean Percentage Change From Baseline at

Enupoint in intention-to-freat i opulation)						
DOSAGE	N	Total-C	LDL-C	HDL-C	TG	Apolipoprotein B
Placebo	47	-1.5	-0.4	-1.9	1.0	0.7
Atorvastatin	140	-31.4	-39.6	2.8	-12.0	-34.0
calcium tablets						
The mean achieved LDL-C value was 130.7 mg/dL (range: 70.0–242.0 mg/dL) in the atorvastatin calcium tablets group compared to 228.5 mg/dL (range: 152.0–385.0 mg/dL) in the placebo group during the 26-week double-blind phase.						

Atorvastatin was also studied in a three year open-label, uncontrolled trial that included 163 patients with HeFH who were 10 years to 15 years old (82 boys and 81 girls). All patients had a clinical diagnosis of HeFH confirmed by genetic analysis (if not already confirmed by family history). Approximately 98% were White, and less than 1% were Black or Asian. Mean LDL-C at baseline was 232 mg/dL. The starting atorvastatin dosage was 10 mg once daily and doses were adjusted to achieve a target of < 130 mg/dL LDL-C. The reductions in LDL-C from baseline were generally consistent across age groups within the trial as well as with previous clinical trials in both adult and pediatric placebo-controlled

16 HOW SUPPLIED/STORAGE AND HANDLING

Atorvastatin calcium tablets are supplied as follows:							
Strength	How Supplied	NDC	Tablet Description				
10 mg of atorvastatin	bottles of 90	60290-039-01	white to off-white, film-coated, oval shaped				
	bottles of 1000	60290-039-02	tablets "ATO" debossed on one side and "10"				
	bottles of 5000	60290-039-03	on other side.				
	bottles of 500	60290-039-04					
20 mg of atorvastatin	bottles of 90	60290-040-01	white to off-white, film-coated, oval shaped				
	bottles of 1000	60290-040-02	tablets "ATO" debossed on one side and "20"				
	bottles of 5000	60290-040-03	on other side.				
	bottles of 500	60290-040-04	1				
40 mg of atorvastatin	bottles of 90	60290-041-01	white to off-white, film-coated, oval shaped				
	bottles of 500	60290-041-02	tablets "ATO" debossed on one side and "40"				
	bottles of 2500	60290-041-03	on other side.				
	bottles of 1000	60290-041-04					
80 mg of atorvastatin	bottles of 90	60290-042-01	white to off-white, film-coated, oval shaped				
	bottles of 500	60290-042-02	tablets "ATO" debossed on one side and "80"				
	bottles of 2000	60290-042-03	on other side.				
	bottles of 1000	60290-042-04					

Store at controlled room temperature 20°C - 25°C (68°F - 77°F)

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

Advise patients that atorvastatin calcium tablets may cause myopathy and rhabdomyolysis. Inform patients that the risk is also increased when taking certain types of medication or consuming large quantities of grapefruit juice and they should discuss all medication, both prescription and over the counter, with their healthcare provider. Instruct patients to promptly report any unexplained muscle pain, tenderness or weakness particularly if accompanied by malaise or fever [see Warnings and Precautions (5.1), Drug Interactions (7.1)].

Inform patients that atorvastatin calcium tablets may cause liver enzyme elevations and possibly liver failure. Advise patients to promptly report fatigue, anorexia, right upper abdominal discomfort, dark urine or jaundice [see Warnings

Increases in HbA1c and Fasting Serum Glucose Levels

Inform patients that increases in HbA1c and fasting serum glucose levels may occur with atorvastatin calcium tablets. Encourage patients to optimize lifestyle measures, including regular exercise, maintaining a healthy body weight, and making healthy food choices [see Warnings and Precautions (5.4)].

Advise pregnant patients and patients who can become pregnant of the potential risk to a fetus. Advise patients to inform their healthcare provider of a known or suspected pregnancy to discuss if atorvastatin calcium tablets should be discontinued [see Use in Specific Populations (8.1)]. Advise patients that breastfeeding is not recommended during treatment with atorvastatin calcium tablets [see Use in

Specific Populations (8.2)]. This product's labeling may have been updated. For the most recent prescribing information, please visit

Umedica Laboratories Pvt. Ltd Plot No. 221 and 221/1, GIDC, IInd Phase

Vapi, Gujarat 396195, INDIA (IND) October 2023; V-06

Hepatic Dysfunction

PATIENT INFORMATION (Atorvastatin Calcium (a tor "va stat' in kal' see um) Tablets USP, for oral use)

Atorvastatin calcium tablets is a prescription medicine that contains a cholesterol lowering medicine (statin) called atorvastatin. Atorvastatin calcium tablets is used: to reduce the risk of:

- o heart attack, stroke, certain types of heart surgery and chest pain in adults who do not have heart disease but have other multiple risk factors for heart disease
- o heart attack and stroke in adults with type 2 diabetes mellitus who do not have heart disease but have other
- o heart attack that does not cause death, stroke, certain types of heart surgery, hospitalization for congestive heart failure, and chest pain in adults with heart dise along with diet to reduce low density lipoprotein cholesterol (LDL-C) or bad cholesterol:
- o in adults with primary hyperlipidemia. o in adults and children aged 10 years and older with heterozygous familial hypercholesterolemia (HeFH). This is an inherited condition that causes high levels of bad cholesterol.
- along with other cholesterol lowering treatments or alone if such treatments are unavailable in adults and children aged 10 years and older with homozygous familial hypercholesterolemia (HoFH). This is an inherited condition that causes high levels of bad cholesterol.

o primary dysbetalipoproteinemia (an inherited condition that causes high levels of cholesterol and fat).

- It is not known if a torvastatin calcium tablets is safe and effective in children younger than 10 years of age with HeFH or HoFH or in children with other types of hyperlipidemias (other than HeFH or HoFH).
- Do not take atorvastatin calcium tablets if you: · have liver problems (acute liver failure or decompensated cirrhosis)

along with diet for the treatment of adults with:

o hypertriglyceridemia.

- are allergic to atorvastatin or any of the ingredients in atorvastatin calcium tablets. Stop using atorvastatin calcium tablets and get medical help right away if you have symptoms of a serious allergic reaction including:
- swelling of your face, lips, tongue or throat o problems breathing or swallowing
- fainting or feeling dizzy o very rapid heartbeat
- o flu-like symptoms including fever, sore throat, cough, tiredness, and ioint pain
- $\underline{\textbf{Before you take Atorva statin calcium tablets, tell your doctor about all of your medical conditions, including if you:}$
- · drink more than 2 glasses of alcohol daily

- are pregnant or plan to become pregnant. Atorvastatin calcium tablets may harm your unborn baby. If you become
 pregnant, stop taking Atorvastatin calcium tablets and call your doctor right away.
- are breastfeeding or plan to breastfeed. You and your doctor should decide if you will take Atorvastatin calcium tablets or breastfeed. You should not do both. Talk to your doctor about the best way to feed your baby if you take Atorvastatin calcium tablets.

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Atoryastatin calcium tablets and certain other medicines can increase the risk of muscle problems or other side effects. Especially tell your doctor if you take medicines for:

- vour immune system (cyclosporine) cholesterol (gemfibrozil)
- $\bullet \quad \text{infections (erythromycin, clarithromycin, itraconazole, ketoconazole, posaconazole, and voriconazole)}\\$ birth control pills
- · heart failure (digoxin)
- · gout (colchicine) niacin
- fibrates
- · viruses that treat HIV, AIDS, or hepatitis C (anti-virals) o tipranavir plus ritonavir o glecaprevir plus pibrentasvii
- ledipasvir plus sofosbuvi o saguinavir plus ritonavir o darunavir plus ritonavi o fosamprenavii o fosamprenavir plus ritonavir

Take atorvastatin calcium tablets exactly as your doctor tells you to take it.

What are the possible side effects of atorvastatin calcium tablets?

o elbasvir plus grazoprevir o letermovir

Ask your doctor or pharmacist for a list of medicines if you are not sure. Know all the medicines you take. Keep a list of them to show your doctor and pharmacist when you get a new medicine. How should I take atorvastatin calcium tablets?

• Do not change your dose or stop atorvastatin calcium tablets without talking to your doctor. Your doctor may do blood tests to check your cholesterol levels during your treatment with atorvastatin calcium tablets. Your dose of atorvastatin calcium tablets may be changed based on these blood test results.

Take atorvastatin calcium tablets each day at any time of day. Atorvastatin calcium tablets can be taken with or Your doctor may start you on a cholesterol lowering diet before giving you atorvastatin calcium tablets. stay on this low-fat diet when you take atorvastatin calcium tablets

If you miss a dose of atorvastatin calcium tablets, take it as soon as you remember. Do not take atorvastatin calcium tablets if it has been more than 12 hours since you missed your last dose. Wait and take the next dose at your regular time. Do not take 2 doses of atorvastatin calcium tablets at the same time. If you take too much atorvastatin calcium tablets or everdose, call your doctor or poison control center at 1-800-222-1222 or go to the

nearest emergency room right away. What should I avoid while taking atorvastatin calcium tablets? . Avoid drinking more than 1.2 liters of grapefruit juice each day.

Atorvastatin calcium tablets may cause serious side effects including: Muscle pain, tenderness and weakness (myopathy). Muscle problems, including muscle breakdown, can be serious in some people and, rarely, cause kidney damage that can lead to death Tell your doctor right away if you have:

o unexplained muscle pain, tenderness, or weakness, especially if you also have a fever or feel more tired than usual while you take atorvastatin calcium tablets. o muscle problems that do not go away after your doctor has told you to stop taking atorvastatin calcium tablets. Your doctor may do further tests to diagnose the cause of your muscle problems.

Your chances of getting muscle problems are higher if you: o are taking certain other medicines while you take atorvastatin calcium tablets o drink large amounts of grapefruit juice o are 65 years of age or older

- o have kidney problems o are taking higher doses of atorvastatin calcium tablets
- **Liver problems.** Your doctor should do blood tests to check your liver before you start taking atorvastatin calcium tablets and if you have symptoms of liver problems while you take atorvastatin calcium tablets. Call your doctor right away if you have the following symptoms of liver problems:
- o nausea or vomiting
- o loss of appetite upper belly pair o dark amber colored urine

o have thyroid problems (hypothyroidism) that are not controlled

- **Increase in blood sugar level.** Your blood sugar level may increase while you are taking atorvastatin calcium tablets. Exercise regularly and make healthy food choices to maintain healthy body weight. The most common side effects of atorvastatin calcium tablets include:
- o nasal congestion, sore throat, runny nose o muscle and joint pain o diarrhea o pain in extremity o urinary tract infectior o upset stomach o nausea o musculoskeletal pain
- o trouble sleeping o throat pain Talk to your doctor or pharmacist if you have side effects that bother you or that will not go away. These are not all the
- effects to FDA at 1-800-FDA-1088

 Store atorvastatin calcium tablets at room temperature between 68° F to 77°F (20°C to 25°C). . Do not keep medicine that is out of date or that you no longer need

Keep atorvastatin calcium tablets and all medicines out of the reach of childre neral Information About the safe and effective use of atorvastatin calcium table Medicines are sometimes prescribed for conditions purposes other than those listed in a Patient Information leaflet. Do not use atorvastatin calcium tablets for a condition for which it was not prescribed. Do not give atorvastatin calcium tablets to other people, even if they have the same symptoms you have. It may harm them. If you would like more information about atorvastatin calcium tablets, talk with your doctor. You can ask your pharmacist or doctor for

information about atorvastatin calcium tablets that is written for health professionals What are the Ingredients in atorvastatin calcium tablets? Active Ingredient: atorvastatin calcium

Inactive Ingredients: Croscarmellose sodium, NF; Hydroxy propyl cellulose, NF, Lactose monohydrate, NF; Magnesium stearate, NF; Microcrystalline cellulose, NF; Polysorbate 80, NF; Precipitated calcium carbonate, NF; Opadry Complete film coating system YS-1-7040 White (Hypromellose, Macrogol, Talc and Titanium dioxide).

Patient Information available at www.umedicalabs.com or call 1-855-288-577

Manufactured by:

Plot No. 221 and 221/1. GIDC. Und Phase. Vapi, Gujarat 396195, INDIA (IND)

October 2023; V-06

400.00 mm